

(FILE 'HOME' ENTERED AT 11:18:16 ON 16 APR 2003)

FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED
AT 11:18:50 ON 16 APR 2003

E RUVKUN G?/AU

L1 76 S E4
L2 69 DUP REM L1 (7 DUPLICATES REMOVED)
L3 11 S L2 AND DAF-16
L4 11 SORT L3 PY
L5 254 S DAF-16
L6 117 DUP REM L5 (137 DUPLICATES REMOVED)
L7 9 S L6 AND PY<=1997
L8 9 SORT L7 PY
L9 80 S L6 AND (EXPRES? OR MODULA? OR INCREAS? OR DECREAS?)
L10 80 FOCUS L9 1-

=> d an ti so au ab l10 2 3 5 8

L10 ANSWER 2 OF 80 MEDLINE

AN 2001699838 MEDLINE

TI Regulation of *C. elegans* **DAF-16** and its human ortholog
FKHRL1 by the *daf-2* insulin-like signaling pathway.

SO CURRENT BIOLOGY, (2001 Dec 11) 11 (24) 1950-7.

Journal code: 9107782. ISSN: 0960-9822.

AU Lee R Y; Hench J; Ruvkun G

AB *C. elegans* insulin-like signaling regulates metabolism, development, and life span. This signaling pathway negatively regulates the activity of the forkhead transcription factor **DAF-16**. *daf-16* encodes multiple isoforms that are expressed in distinct tissue types and are probable orthologs of human FKHRL1, FKHR, and AFX. We show that human FKHRL1 can partially replace **DAF-16**, proving the orthology. In mammalian cells, insulin and insulin-like growth factor signaling activate AKT/PKB kinase to negatively regulate the nuclear localization of **DAF-16** homologs (reviewed in). We show that the absence of AKT consensus sites on **DAF-16** is sufficient to cause dauer arrest in *daf-2*(+) animals, proving that *daf-16* is the major output of insulin signaling in *C. elegans*. FKHR, FKHRL1, and AFX may similarly be the major outputs of mammalian insulin signaling. *daf-2* insulin signaling, via AKT kinases, negatively regulates **DAF-16** by controlling its nuclear localization. Surprisingly, we find that *daf-7* TGF-beta signaling also regulates **DAF-16** nuclear localization specifically at the time when the animal makes the commitment between diapause and reproductive development. *daf-16* function is supported by the combined action of two distinct promoter/enhancer elements, whereas the coding sequences of two major **DAF-16** isoforms are interchangeable. Together, these observations suggest that the combined effects of transcriptional and posttranslational regulation of *daf-16* transduce insulin-like signals in *C. elegans* and perhaps more generally.

L10 ANSWER 3 OF 80 MEDLINE

AN 1998013175 MEDLINE

TI The Fork head transcription factor **DAF-16** transduces
insulin-like metabolic and longevity signals in *C. elegans*.

SO NATURE, (1997 Oct 30) 389 (6654) 994-9.

Journal code: 0410462. ISSN: 0028-0836.

AU Ogg S; Paradis S; Gottlieb S; Patterson G I; Lee L; Tissenbaum H A; Ruvkun G

AB In mammals, insulin signalling regulates glucose transport together with the **expression** and activity of various metabolic enzymes. In the nematode *Caenorhabditis elegans*, a related pathway regulates metabolism, development and longevity. Wild-type animals enter the developmentally arrested dauer stage in response to high levels of a secreted pheromone, accumulating large amounts of fat in their intestines and hypodermis. Mutants in *DAF-2* (a homologue of the mammalian insulin receptor) and *AGE-1* (a homologue of the catalytic subunit of mammalian phosphatidylinositol 3-OH kinase) arrest development at the dauer stage. Moreover, animals bearing weak or temperature-sensitive mutations in *daf-2*

and age-1 can develop reproductively, but nevertheless show **increased** energy storage and longevity. Here we show that null mutations in **daf-16** suppress the effects of mutations in **daf-2** or **age-1**; lack of **daf-16** bypasses the need for this insulin receptor-like signalling pathway. The principal role of DAF-2/AGE-1 signalling is thus to antagonize **DAF-16**. **daf-16** is widely **expressed** and encodes three members of the Fork head family of transcription factors. The DAF-2 pathway acts synergistically with the pathway activated by a nematode TGF-beta-type signal, DAF-7, suggesting that **DAF-16** cooperates with nematode SMAD proteins in regulating the transcription of key metabolic and developmental control genes. The probable human orthologues of **DAF-16**, FKHR and AFX, may also act downstream of insulin signalling and cooperate with TGF-beta effectors in mediating metabolic regulation. These genes may be dysregulated in diabetes.

L10 ANSWER 5 OF 80 MEDLINE
 AN 2001698389 MEDLINE
 TI **DAF-16**-dependent and independent **expression** targets of DAF-2 insulin receptor-like pathway in *Caenorhabditis elegans* include FKBP's.
 SO JOURNAL OF MOLECULAR BIOLOGY, (2001 Dec 14) 314 (5) 1017-28.
 Journal code: 2985088R. ISSN: 0022-2836.
 AU Yu H; Larsen P L
 AB The daf-2 insulin-like receptor pathway regulates development and life-span in *Caenorhabditis elegans*. Reduced DAF-2 signaling leads to changes in downstream targets via the **daf-16** gene, a fork-head transcription factor which is regulated by DAF-2, and results in extended life-span. Here, we describe the first identification of genes whose **expression** is controlled by the DAF-2 signaling cascade. **dao-1**, **dao-2**, **dao-3**, **dao-4**, **dao-8** and **dao-9** are down-regulated in **daf-2** mutant adults compared to wild-type adults, whereas **dao-5**, **dao-6** and **dao-7** are up-regulated. The latter genes are negatively regulated by DAF-2 signaling and positively regulated by **DAF-16**. Positive regulation by DAF-2 on **dao-1**, **dao-4** and **dao-8** was mediated by **DAF-16**, whereas **daf-16** mediates only part of DAF-2 signaling for **dao-2** and **dao-9**. Regulation by DAF-2 is most likely **DAF-16** independent for **dao-3** and **hsp-90**. RNA levels of **dao-5** and **dao-6** showed elevated **expression** in **daf-2** adults, as well as being strongly **expressed** in dauer larvae. In contrast, **hsp-90** transcript levels are low in **daf-2** mutant adults though they are enriched in dauer larvae, indicating overlapping but not identical mechanisms of efficient life maintenance in stress-resistant dauer larvae and long-lived **daf-2** mutant adults. **dao-1**, **dao-8** and **dao-9** are homologs of the FK506 binding proteins that interact with the mammalian insulin pathway. **dao-3** encodes a putative methylenetetrahydrofolate dehydrogenase. **DAO-5** shows 33 % identity with human nucleolar phosphoprotein P130. **dao-7** is similar to the mammalian ZFP36 protein. Distinct regulatory patterns of **dao** genes implicate their diverse positions within the signaling network of DAF-2 pathway, and suggest they have unique contributions to development, metabolism and longevity.
 Copyright 2001 Academic Press.

L10 ANSWER 8 OF 80 MEDLINE
 AN 1998028757 MEDLINE
 TI **daf-16**: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*.
 SO SCIENCE, (1997 Nov 14) 278 (5341) 1319-22.
 Journal code: 0404511. ISSN: 0036-8075.
 AU Lin K; Dorman J B; Rodan A; Kenyon C
 AB The wild-type *Caenorhabditis elegans* nematode ages rapidly, undergoing development, senescence, and death in less than 3 weeks. In contrast, mutants with reduced activity of the gene **daf-2**, a homolog of the insulin and insulin-like growth factor receptors, age more slowly than normal and live more than twice as long. These mutants are active and fully fertile and have normal metabolic rates. The life-span extension caused by **daf-2** mutations requires the activity of the gene **daf-16**. **daf-16** appears to play a unique role in life-span

regulation and encodes a member of the hepatocyte nuclear factor 3 (HNF-3)/forkhead family of transcriptional regulators. In humans, insulin down-regulates the **expression** of certain genes by antagonizing the activity of HNF-3, raising the possibility that aspects of this regulatory system have been conserved.

=>

(FILE 'HOME' ENTERED AT 11:18:16 ON 16 APR 2003)

FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED
AT 11:18:50 ON 16 APR 2003

E RUVKUN G?/AU

L1 76 S E4
L2 69 DUP REM L1 (7 DUPLICATES REMOVED)
L3 11 S L2 AND DAF-16
L4 11 SORT L3 PY
L5 254 S DAF-16
L6 117 DUP REM L5 (137 DUPLICATES REMOVED)
L7 9 S L6 AND PY<=1997
L8 9 SORT L7 PY

=> d an ti so au ab pi l8 1-9

L8 ANSWER 1 OF 9 MEDLINE
AN 92120509 MEDLINE
TI Genetic analysis of chemosensory control of dauer formation in
Caenorhabditis elegans.
SO GENETICS, (1992 Jan) 130 (1) 105-23.
Journal code: 0374636. ISSN: 0016-6731.
AU Vowels J J; Thomas J H
AB Dauer larva formation in Caenorhabditis elegans is controlled by
chemosensory cells that respond to environmental cues. Genetic
interactions among mutations in 23 genes that affect dauer larva formation
were investigated. Mutations in seven genes that cause constitutive dauer
formation, and mutations in 16 genes that either block dauer formation or
result in the formation of abnormal dauers, were analyzed. Double mutants
between dauer-constitutive and dauer-defective mutations were constructed
and characterized for their capacity to form dauer larvae. Many of the
genes could be interpreted to lie in a simple linear epistasis pathway.
Three genes, *daf-16*, *daf-18* and *daf-20*, may affect
downstream steps in a branched part of the pathway. Three other genes,
daf-2, *daf-3* and *daf-5*, displayed partial or complex epistasis
interactions that were difficult to interpret as part of a simple linear
pathway. Dauer-defective mutations in nine genes cause structurally
defective chemosensory cilia, thereby blocking chemosensation. Mutations
in all nine of these genes appear to fall at a single step in the
epistasis pathway. Dauer-constitutive mutations in one gene, *daf-11*, were
strongly suppressed for dauer formation by mutations in the nine
cilium-structure genes. Mutations in the other six dauer-constitutive
genes caused dauer formation despite the absence of functional
chemosensory endings. These results suggest that *daf-11* is directly
involved in chemosensory transduction essential for dauer formation, while
the other *Daf-c* genes play roles downstream of the chemosensory step.

L8 ANSWER 2 OF 9 MEDLINE
AN 94067343 MEDLINE
TI A C. elegans mutant that lives twice as long as wild type.
SO NATURE, (1993 Dec 2) 366 (6454) 461-4.
Journal code: 0410462. ISSN: 0028-0836.
AU Kenyon C; Chang J; Gensch E; Rudner A; Tabtiang R
AB We have found that mutations in the gene *daf-2* can cause fertile, active,
adult Caenorhabditis elegans hermaphrodites to live more than twice as
long as wild type. This lifespan extension, the largest yet reported in
any organism, requires the activity of a second gene, *daf-16*. Both genes also regulate formation of the dauer larva, a
developmentally arrested larval form that is induced by crowding and
starvation and is very long-lived. Our findings raise the possibility
that the longevity of the dauer is not simply a consequence of its
arrested growth, but instead results from a regulated lifespan extension
mechanism that can be uncoupled from other aspects of dauer formation.
daf-2 and *daf-16* provide entry points into
understanding how lifespan can be extended.

L8 ANSWER 3 OF 9 MEDLINE
AN 94333774 MEDLINE
TI *daf-2*, *daf-16* and *daf-23*: genetically interacting
genes controlling Dauer formation in Caenorhabditis elegans.

SO GENETICS, (1994 May) 137 (1) 107-20.
Journal code: 0374636. ISSN: 0016-6731.

AU Gottlieb S; Ruvkun G
AB Under conditions of high population density and low food, *Caenorhabditis elegans* forms an alternative third larval stage, called the dauer stage, which is resistant to desiccation and harsh environments. Genetic analysis of some dauer constitutive (Daf-c) and dauer defective (Daf-d) mutants has revealed a complex pathway that is likely to function in particular neurons and/or responding tissues. Here we analyze the genetic interactions between three genes which comprise a branch of the dauer formation pathway that acts in parallel to or downstream of the other branches of the pathway, the Daf-c genes *daf-2* and *daf-23* and the Daf-d gene *daf-16*. Unlike mutations in other Daf-c genes, mutations in both *daf-2* and *daf-23* cause non-conditional arrest at the dauer stage. Our epistasis analysis suggests that *daf-2* and *daf-23* are functioning at a similar point in the dauer pathway. First, mutations in *daf-2* and *daf-23* are epistatic to mutations in the same set of Daf-d genes. Second, *daf-2* and *daf-23* mutants are suppressed by mutations in *daf-16*. Mutations in *daf-16* do not suppress any of the other Daf-c mutants as efficiently as they suppress *daf-2* and *daf-23* mutants. Third, double mutants between either *daf-2* or *daf-23* and several other *daf-d* mutants exhibit an unusual interaction. Based on these results, we present a model for the function of *daf-2*, *daf-23* and *daf-16* in dauer formation.

L8 ANSWER 4 OF 9 MEDLINE
AN 96170778 MEDLINE
TI The age-1 and *daf-2* genes function in a common pathway to control the lifespan of *Caenorhabditis elegans*.
SO GENETICS, (1995 Dec) 141 (4) 1399-406.
Journal code: 0374636. ISSN: 0016-6731.

AU Dorman J B; Albinder B; Shroyer T; Kenyon C
AB Recessive mutations in two genes, *daf-2* and *age-1*, extend the lifespan of *Caenorhabditis elegans* significantly. The *daf-2* gene also regulates formation of an alternative developmental state called the dauer. Here we asked whether these two genes function in the same or different lifespan pathways. We found that the longevity of both *age-1* and *daf-2* mutants requires the activities of the same two genes, *daf-16* and *daf-18*. In addition, the *daf-2*(*el370*); *age-1*(*hx546*) double mutant did not live significantly longer than the *daf-2* single mutant. We also found that, like *daf-2* mutations, the *age-1*(*hx546*) mutation affects certain aspects of dauer formation. These findings suggest that *age-1* and *daf-2* mutations do act in the same lifespan pathway and extend lifespan by triggering similar if not identical processes.

L8 ANSWER 5 OF 9 MEDLINE
AN 95309673 MEDLINE
TI Genes that regulate both development and longevity in *Caenorhabditis elegans*.
SO GENETICS, (1995 Apr) 139 (4) 1567-83.
Journal code: 0374636. ISSN: 0016-6731.

AU Larsen P L; Albert P S; Riddle D L
AB The nematode *Caenorhabditis elegans* responds to conditions of overcrowding and limited food by arresting development as a dauer larva. Genetic analysis of mutations that alter dauer larva formation (*daf* mutations) is presented along with an updated genetic pathway for dauer vs. nondauer development. Mutations in the *daf-2* and *daf-23* genes double adult life span, whereas mutations in four other dauer-constitutive genes positioned in a separate branch of this pathway (*daf-1*, *daf-4*, *daf-7* and *daf-8*) do not. The increased life spans are suppressed completely by a *daf-16* mutation and partially in a *daf-2*; *daf-18* double mutant. A genetic pathway for determination of adult life span is presented based on the same strains and growth conditions used to characterize *Daf* phenotypes. Both dauer larva formation and adult life span are affected in *daf-2*; *daf-12* double mutants in an allele-specific manner. Mutations in *daf-12* do not extend adult life span, but certain combinations of *daf-2* and *daf-12* mutant alleles nearly quadruple it. This synergistic effect, which does not equivalently extend the fertile period, is the largest genetic extension of life span yet observed in a metazoan.

L8 ANSWER 6 OF 9 MEDLINE
 AN 96400918 MEDLINE
 TI A genetic pathway conferring life extension and resistance to UV stress in *Caenorhabditis elegans*.
 SO GENETICS, (1996 Jul) 143 (3) 1207-18.
 Journal code: 0374636. ISSN: 0016-6731.
 AU Murakami S; Johnson T E
 AB A variety of mechanisms have been proposed to explain the extension of adult life span (Age) seen in several mutants in *Caenorhabditis elegans* (age-1: an altered aging rate; daf-2 and daf-23: activation of a dauer-specific longevity program; spe-26: reduced fertility; clk-1: an altered biological clock). Using an assay for ultraviolet (UV) resistance in young adult hermaphrodites (survival after UV irradiation), we observed that all these Age mutants show increased resistance to UV. Moreover, mutations in **daf-16** suppressed the UV resistance as well as the increased longevity of all the Age mutants. In contrast to the multiple mechanisms initially proposed, these results suggest that a single, **daf-16**-dependent pathway, specifies both extended life span and increased UV resistance. The mutations in **daf-16** did not alter the reduced fertility of spe-26 and interestingly a **daf-16** mutant is more fertile than wild type. We propose that life span and some aspects of stress resistance are jointly negatively regulated by a set of gerontogenes (genes whose alteration causes life extension) in *C. elegans*.

L8 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2003 ACS
 AN 1996:303001 CAPLUS
 DN 125:5796
 TI Determination of life-span in *Caenorhabditis elegans* by four Clock genes
 SO Science (Washington, D. C.) (1996), 272(5264), 1010-1013
 CODEN: SCIEAS; ISSN: 0036-8075
 AU Lakowski, Bernard; Hekimi, Siegfried
 AB The nematode worm *Caenorhabditis elegans* is a model system for the study of the genetic basis of aging. Maternal-effect mutations in four genes-clk-1, clk-2, clk-3, and gro-1-interact genetically to det. both the duration of development and life-span. Anal. of the phenotypes of these mutants suggests the existence of a general physiol. clock in the worm. Mutations in certain genes involved in dauer formation (an alternative larval stage induced by adverse conditions in which development is arrested) can also extend life-span, but the life extension of Clock mutants appears to be independent of these genes. The daf-2(e1370) clk-1(e2519) worms, which carry life-span-extending mutations from two different pathways, live nearly five times as long as wild-type worms.

L8 ANSWER 8 OF 9 MEDLINE
 AN 1998028757 MEDLINE
 TI **daf-16**: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*.
 SO SCIENCE, (1997 Nov 14) 278 (5341) 1319-22.
 Journal code: 0404511. ISSN: 0036-8075.
 AU Lin K; Dorman J B; Rodan A; Kenyon C
 AB The wild-type *Caenorhabditis elegans* nematode ages rapidly, undergoing development, senescence, and death in less than 3 weeks. In contrast, mutants with reduced activity of the gene daf-2, a homolog of the insulin and insulin-like growth factor receptors, age more slowly than normal and live more than twice as long. These mutants are active and fully fertile and have normal metabolic rates. The life-span extension caused by daf-2 mutations requires the activity of the gene **daf-16**. **daf-16** appears to play a unique role in life-span regulation and encodes a member of the hepatocyte nuclear factor 3 (HNF-3)/forkhead family of transcriptional regulators. In humans, insulin down-regulates the expression of certain genes by antagonizing the activity of HNF-3, raising the possibility that aspects of this regulatory system have been conserved.

L8 ANSWER 9 OF 9 MEDLINE
 AN 1998013175 MEDLINE
 TI The Fork head transcription factor **DAF-16** transduces insulin-like metabolic and longevity signals in *C. elegans*.
 SO NATURE, (1997 Oct 30) 389 (6654) 994-9.

Journal code: 0410462. ISSN: 0028-0836.

AU Ogg S; Paradis S; Gottlieb S; Patterson G I; Lee L; Tissenbaum H A; Ruvkun G

AB In mammals, insulin signalling regulates glucose transport together with the expression and activity of various metabolic enzymes. In the nematode *Caenorhabditis elegans*, a related pathway regulates metabolism, development and longevity. Wild-type animals enter the developmentally arrested dauer stage in response to high levels of a secreted pheromone, accumulating large amounts of fat in their intestines and hypodermis. Mutants in DAF-2 (a homologue of the mammalian insulin receptor) and AGE-1 (a homologue of the catalytic subunit of mammalian phosphatidylinositol 3-OH kinase) arrest development at the dauer stage. Moreover, animals bearing weak or temperature-sensitive mutations in *daf-2* and *age-1* can develop reproductively, but nevertheless show increased energy storage and longevity. Here we show that null mutations in **daf-16** suppress the effects of mutations in *daf-2* or *age-1*; lack of **daf-16** bypasses the need for this insulin receptor-like signalling pathway. The principal role of DAF-2/AGE-1 signalling is thus to antagonize **DAF-16**. **daf-16** is widely expressed and encodes three members of the Fork head family of transcription factors. The DAF-2 pathway acts synergistically with the pathway activated by a nematode TGF-beta-type signal, DAF-7, suggesting that **DAF-16** cooperates with nematode SMAD proteins in regulating the transcription of key metabolic and developmental control genes. The probable human orthologues of **DAF-16**, FKHR and AFX, may also act downstream of insulin signalling and cooperate with TGF-beta effectors in mediating metabolic regulation. These genes may be dysregulated in diabetes.

=>

L4 ANSWER 2 OF 11 CAPLUS COPYRIGHT 2003 ACS
 AN 1997:702845 CAPLUS
 DN 128:20747
 TI The Fork head transcription factor **DAF-16** transduces insulin-like metabolic and longevity signals in *C. elegans*
 SO Nature (London) (1997), 389(6654), 994-999
 CODEN: NATUAS; ISSN: 0028-0836
 AU Ogg, Scott; Paradis, Suzanne; Gottlieb, Shoshanna; Patterson, Garth I.; Lee, Linda; Tissenbaum, Heidi A.; **Ruvkun, Gary**
 AB In mammals, insulin signaling regulates glucose transport together with the expression and activity of various metabolic enzymes. In the nematode *Caenorhabditis elegans*, a related pathway regulates metab., development and longevity. Wild-type animals enter the developmentally arrested dauer stage in response to high levels of a secreted pheromone, accumulating large amts. of fat in their intestines and hypodermis. Mutants in **DAF-2** (a homolog of the mammalian insulin receptor) and **AGE-1** (a homolog of the catalytic subunit of mammalian phosphatidylinositol 3-OH kinase) arrest development at the dauer stage. Moreover, animals bearing weak or temp.-sensitive mutations in **daf-2** and **age-1** can develop reproductively, but nevertheless show increased energy storage and longevity. Null mutations in **daf-16** suppress the effects of mutations in **daf-2** or **age-1**; lack of **daf-16** bypasses the need for this insulin receptor-like signaling pathway. **DAF-16** is widely expressed and encodes three members of the Fork head family of transcription factors. The **DAF-2** pathway acts synergistically with the pathway activated by a nematode TGF- β -type signal, **DAF-7**, suggesting that **DAF-16** cooperates with nematode SMAD proteins in regulating the transcription of key metabolic and developmental control genes. The probable human orthologs of **DAF-16**, **FKHR** and **AFX**, may also act downstream of insulin signaling and cooperate with TGF- β effectors in mediating metabolic regulation. These genes may be dysregulated in diabetes.

L4 ANSWER 3 OF 11 CAPLUS COPYRIGHT 2003 ACS
 AN 1999:37224 CAPLUS
 DN 130:220650
 TI The *C. elegans* PTEN homolog, **DAF-18**, acts in the insulin receptor-like metabolic signaling pathway
 SO Molecular Cell (1998), 2(6), 887-893
 CODEN: MOCEFL; ISSN: 1097-2765
 AU Ogg, Scott; **Ruvkun, Gary**
 AB An insulin-like signaling pathway, from the **DAF-2** receptor, the **AGE-1** phosphoinositide 3-kinase, and the **AKT-1/AKT-2** serine/threonine kinases to the **DAF-16** Fork head transcription factor, regulates the metab., development, and life span of *Caenorhabditis elegans*. Inhibition of **daf-18** gene activity bypasses the normal requirement for **AGE-1** and partially bypasses the need for **DAF-2** signaling. The suppression of **age-1** mutations by a **daf-18** mutation depends on **AKT-1/AKT-2** signaling, showing that **DAF-18** acts between **AGE-1** and the **AKT** input to **DAF-16** transcriptional regulation. **Daf-18** encodes a homolog of the human tumor suppressor PTEN (MMAC1/TEP1), which has 3-phosphatase activity toward phosphatidylinositol 3,4,5-trisphosphate (PIP3). **DAF-18** PTEN may normally limit **AKT-1** and **AKT-2** activation by decreasing PIP3 levels. The action of **daf-18** in this metabolic control pathway suggests that mammalian PTEN may modulate insulin signaling and may be variant in diabetic pedigrees.

L4 ANSWER 4 OF 11 CAPLUS COPYRIGHT 2003 ACS
 AN 1998:761816 CAPLUS
 DN 130:29188
 TI Therapeutic and diagnostic tools for impaired glucose tolerance conditions based on the dauer polypeptides and genes of *Caenorhabditis elegans*
 SO PCT Int. Appl., 202 pp.
 CODEN: PIXXD2
 IN **Ruvkun, Gary**; Kimura, Koutarou; Patterson, Garth; Ogg, Scott; Paradis, Suzanne; Tissenbaum, Heidi; Morris, Jason; Koweeek, Allison; Pierce, Sarah
 AB Disclosed herein are novel genes and methods for the screening of therapeutics useful for treating impaired glucose tolerance conditions, as

well as diagnostics and therapeutic compns. for identifying or treating such conditions. The *Caenorhabditis elegans* metabolic regulatory genes *daf-2* and *age-1* encode homologs of the mammalian insulin receptor/phosphoinositide 3-kinase signaling pathway proteins, resp. In addn., the **DAF-16** forkhead protein represents the major transcriptional output of this insulin signaling pathway. Dysregulation of the **DAF-16** transcription factor in the absence of insulin signaling leads to metabolic defects; inactivation of **DAF-16** reverses the metabolic defects caused by lack of insulin signaling in *C. elegans*. Finally, the *C. elegans* *daf-7*, *da-1*, *daf-4*, *daf-8*, *daf-14*, and *daf-3* genes encode neuroendocrine/target tissue transforming growth factor- β . type signal transduction mols. that genetically interact with the insulin signaling pathway. Metabolic defects cause by lack of neuroendocrine TGF- β . signals can be reversed by inactivation of the **DAF-3** transcription factor. The *C. elegans* *daf* genes are excellent candidate genes and proteins for human disease assocd. with glucose intolerance, e.g., diabetes, obesity, and atherosclerosis. The human homologs of these *daf* genes and proteins mediate insulin signaling in normal people and may be defective or mis-regulated in diabetics. Moreover, there are at least 2 classes of type II diabetics: those with defects in the TGF- β . signaling genes, and those with defects in insulin signaling genes. Exemplary sequences and functional characteristics are provided for the *C. elegans* *daf* homologs of the human genes: *daf-2*, *daf-3* (3 differentially spliced isoforms), **daf-16** (2 differentially spliced isoforms), *age-1*, and *pdk-1* (two spliced isoforms).

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 9851351	A1	19981119	WO 1998-US10080	19980515
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
US 6225120	B1	20010501	US 1997-857076	19970515
AU 9874941	A1	19981208	AU 1998-74941	19980515
AU 752962	B2	20021003		
EP 1019092	A1	20000719	EP 1998-922382	19980515
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
JP 2002511747	T2	20020416	JP 1998-549639	19980515
US 2001029617	A1	20011011	US 1998-205658	19981203
US 2002037585	A1	20020328	US 2001-844353	20010427

L4 ANSWER 5 OF 11 CAPLUS COPYRIGHT 2003 ACS

AN 1998:577164 CAPLUS

DN 129:287966

TI *Caenorhabditis elegans* Akt/PKB transduces insulin receptor-like signals from AGE-1 PI3 kinase to the **DAF-16** transcription factor

SO Genes & Development (1998), 12(16), 2488-2498
CODEN: GEDEEP; ISSN: 0890-9369

AU Paradis, Suzanne; Ruvkun, Gary

AB A neurosecretory pathway regulates a reversible developmental arrest and metabolic shift at the *Caenorhabditis elegans* dauer larval stage. Defects in an insulin-like signaling pathway cause arrest at the dauer stage. We show here that two *C. elegans* Akt/PKB homologs, *akt-1* and *akt-2*, transduce insulin receptor-like signals that inhibit dauer arrest and that AKT-1 and AKT-2 signaling are indispensable for insulin receptor-like signaling in *C. elegans*. A loss-of-function mutation in the Fork head transcription factor **DAF-16** relieves the requirement for Akt/PKB signaling, which indicates that AKT-1 and AKT-2 function primarily to antagonize **DAF-16**. This is the first evidence that the major target of Akt/PKB signaling is a transcription factor. An activation mutation in *akt-1*, revealed by a genetic screen, as well as increased dosage of wild-type *akt-1* relieves the requirements for signaling from AGE-1 PI3K, which acts downstream of the **DAF-02**

insulin/IGF-1 receptor homolog. This demonstrates that Akt/PKB activity is not necessarily dependent on AGE-1 PI3K activity. Akt-1 and akt-2 are expressed in overlapping patterns in the nervous system and in tissues that are remodeled during dauer formation.

L4 ANSWER 6 OF 11 CAPLUS COPYRIGHT 2003 ACS
AN 1998:463129 CAPLUS
DN 129:228428
TI An insulin-like signaling pathway affects both longevity and reproduction in *Caenorhabditis elegans*
SO Genetics (1998), 148(2), 703-717
CODEN: GENTAE; ISSN: 0016-6731
AU Tissenbaum, Heidi A.; Ruvkun, Gary
AB Mutations in daf-2 and age-1 cause a dramatic increase in longevity as well as developmental arrest at the dauer diapause stage in *Caenorhabditis elegans*. Daf-2 and age-1 encode components of an insulin-like signaling pathway. Both daf-2 and age-1 act at a similar point in the genetic epistasis pathway for dauer arrest and longevity and regulate the activity of the daf-16 gene. Mutations in daf-16 cause a dauer-defective phenotype and are epistatic to the diapause arrest and life span extension phenotypes of daf-2 and age-1 mutants. Mutations in this pathway also affect fertility and embryonic development. Weak daf-2 alleles, and maternally rescued age-1 alleles that cause life span extension but do not arrest at the dauer stage, also reduce fertility and viability. The authors find that age-1(hx546) has reduced both maternal and zygotic age-1 activity. Daf-16 mutations suppress all of the daf-2 and age-1 phenotypes, including dauer arrest, life span extension, reduced fertility, and viability defects. These data show that insulin signaling, mediated by DAF-2 through the AGE-1 phosphatidylinositol-3-hydroxykinase, regulates reprodn. and embryonic development, as well as dauer diapause and life span, and that DAF-16 transduces these signals. The regulation of fertility, life span, and metab. by an insulin-like signaling pathway is similar to the endocrine regulation of metab. and fertility by mammalian insulin signaling.

L4 ANSWER 7 OF 11 CAPLUS COPYRIGHT 2003 ACS
AN 1999:382232 CAPLUS
DN 131:182560
TI A PDK1 homolog is necessary and sufficient to transduce AGE-1 PI3 kinase signals that regulate diapause in *Caenorhabditis elegans*
SO Genes & Development (1999), 13(11), 1438-1452
CODEN: GEDEEP; ISSN: 0890-9369
AU Paradis, Suzanne; Ailion, Michael; Toker, Alex; Thomas, James H.; Ruvkun, Gary
AB An insulin receptor-like signaling pathway regulates *Caenorhabditis elegans* metab., development, and longevity. Inactivation of the insulin receptor homolog DAF-2, the AGE-1 PI3K, or the AKT-1 and AKT-2 kinases causes a developmental arrest at the dauer stage. A null mutation in the daf-16 Fork head transcription factor alleviates the requirement for signaling through this pathway. We show here that a loss-of-function mutation in pdk-1, the *C. elegans* homolog of the mammalian Akt/PKB kinase PDK1, results in constitutive arrest at the dauer stage and increased life span; these phenotypes are suppressed by a loss of function mutation in daf-16. An activating mutation in pdk-1 or overexpression of wild-type pdk-1 relieves the requirement for AGE-1 PI3K signaling. Therefore, pdk-1 activity is both necessary and sufficient to propagate AGE-1 PI3K signals in the DAF-2 insulin receptor-like signaling pathway. The activating mutation in pdk-1 requires akt-1 and akt-2 gene activity in order to suppress the dauer arrest phenotype of age-1. This indicates that the major function of *C. elegans* PDK1 is to transduce signals from AGE-1 to AKT-1 and AKT-2. The activating pdk-1 mutation is located in a conserved region of the kinase domain; the equiv. amino acid substitution in human PDK1 activates its kinase activity toward mammalian Akt/PKB.

L4 ANSWER 8 OF 11 CAPLUS COPYRIGHT 2003 ACS
AN 2000:664827 CAPLUS
DN 134:158393
TI DAF-16 recruits the CREB-binding protein coactivator

complex to the insulin-like growth factor binding protein 1 promoter in HepG2 cells

SO Proceedings of the National Academy of Sciences of the United States of America (2000), 97(19), 10412-10417
CODEN: PNASA6; ISSN: 0027-8424

AU Nasrin, Nargis; Ogg, Scott; Cahill, Catherine M.; Biggs, William; Nui, Simin; Dore, Justin; Calvo, Dominica; Shi, Yang; **Ruvkun, Gary**; Alexander-Bridges, Maria C.

AB Insulin neg. regulates expression of the insulin-like growth factor binding protein 1 (IGFBP-1) gene by means of an insulin-responsive element (IRE) that also contributes to glucocorticoid stimulation of this gene. We find that the *Caenorhabditis elegans* protein **DAF-16** binds the IGFBP-1.cntdot.IRE with specificity similar to that of the forkhead (FKH) factor(s) that act both to enhance glucocorticoid responsiveness and to mediate the neg. effect of insulin at this site. In HepG2 cells, **DAF-16** and its mammalian homologs, FKHR, FKHL1, and AFX, activate transcription through the IGFBP-1.cntdot.IRE; this effect is inhibited by the viral oncoprotein E1A, but not by mutants of E1A that fail to interact with the coactivator p300/CREB-binding protein (CBP). We show that **DAF-16** and FKHR can interact with both the KIX and E1A/SRC interaction domains of p300/CBP, as well as the steroid receptor coactivator (SRC). A C-terminal deletion mutant of **DAF-16** that is nonfunctional in *C. elegans* fails to bind the KIX domain of CBP, fails to activate transcription through the IGFBP-1.cntdot.IRE, and inhibits activation of the IGFBP-1 promoter by glucocorticoids. Thus, the interaction of **DAF-16** homologs with the KIX domain of CBP is essential to basal and glucocorticoid-stimulated transactivation. Although AFX interacts with the KIX domain of CBP, it does not interact with SRC and does not respond to glucocorticoids or insulin. Thus, we conclude that **DAF-16** and FKHR act as accessory factors to the glucocorticoid response, by recruiting the p300/CBP/SRC coactivator complex to an FKH factor site in the IGFBP-1 promoter, which allows the cell to integrate the effects of glucocorticoids and insulin on genes that carry this site.

L4 ANSWER 9 OF 11 CAPLUS COPYRIGHT 2003 ACS

AN 2000:384548 CAPLUS

DN 133:39116

TI Genes and polypeptides involved in insulin signaling pathways for glucose tolerance, obesity, and longevity and their uses as therapeutic and diagnostic tools

SO PCT Int. Appl., 402 pp.
CODEN: PIXXD2

IN **Ruvkun, Gary**; Ogg, Scott

AB Disclosed herein are novel genes and methods for the screening of therapeutics useful for treating impaired glucose tolerance conditions, as well as diagnostics and therapeutic compns. for identifying or treating such conditions. The *Caenorhabditis elegans* metabolic regulatory genes *daf-2* and *age-1* encode homologs of the mammalian insulin receptor/phosphoinositol 3-kinase signaling pathway proteins, resp. Also, the *C. elegans* PKB kinase and AKT kinase act downstream of these genes, as their mammalian homologs act downstream of insulin signaling. The *C. elegans* PTEN lipid phosphatase homolog, *DAF-18*, acts upstream of AKT in this signaling pathway. Further, the **DAF-16** forkhead protein represents the major transcriptional output of this insulin signaling pathway. Addnl. evidence indicates that the **DAF-16**, *DAF-3*, *DAF-8*, and *DAF-14* transcriptional outputs of converging signaling pathways regulate metab. The congruence between the *C. elegans* and mammalian insulin signaling pathways strongly supports the contention that new genes identified in the *C. elegans* pathway also act in mammalian insulin signaling. Exemplary sequences and functional characteristics of the *C. elegans* *daf* genes and their human homologs are provided.

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000033068	A1	20000608	WO 1999-US28529	19991202

PI W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ,

BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,
 DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,
 CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 US 2001029617 A1 20011011 US 1998-205658 19981203
 EP 1163515 A1 20011219 EP 1999-960641 19991202
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO

L4 ANSWER 10 OF 11 CAPLUS COPYRIGHT 2003 ACS
 AN 2002:2553 CAPLUS
 DN 136:381187
 TI Regulation of *C. elegans* **DAF-16** and its human ortholog
 FKHL1 by the *daf-2* insulin-like signaling pathway
 SO Current Biology (2001), 11(24), 1950-1957
 CODEN: CUBLE2; ISSN: 0960-9822
 AU Lee, Raymond Y. N.; Hench, Jurgen; **Ruvkun, Gary**
 AB *C. elegans* insulin-like signaling regulates metab., development, and life
 span. This signaling pathway neg. regulates the activity of the forkhead
 transcription factor **DAF-16**. **Daf-16**
 encodes multiple isoforms that are expressed in distinct tissue types and
 are probable orthologs of human FKHL1, FKHL, and AFX. The authors show
 that human FKHL1 can partially replace **DAF-16**,
 proving the orthol. In mammalian cells, insulin and insulin-like growth
 factor signaling activate AKT/PKB kinase to neg. regulate the nuclear
 localization of **DAF-16** homologs. The authors show
 that the absence of AKT consensus sites on **DAF-16** is
 sufficient to cause dauer arrest in *daf-2(+)* animals, proving that
daf-16 is the major output of insulin signaling in *C.*
elegans. FKHL, FKHL1, and AFX may similarly be the major outputs of
 mammalian insulin signaling. *Daf-2* insulin signaling, via AKT kinases,
 neg. regulates **DAF-16** by controlling its nuclear
 localization. Surprisingly, the authors find that *daf-7* TGF- β .
 signaling also regulates **DAF-16** nuclear localization
 specifically at the time when the animal makes the commitment between
 diapause and reproductive development. **Daf-16**
 function is supported by the combined action of two distinct
 promoter/enhancer elements, whereas the coding sequences of two major
DAF-16 isoforms are interchangeable. Together, these
 observations suggest that the combined effects of transcriptional and
 posttranslational regulation of *daf-16* transduce
 insulin-like signals in *C. elegans* and perhaps more generally.

L4 ANSWER 11 OF 11 CAPLUS COPYRIGHT 2003 ACS
 AN 2001:318557 CAPLUS
 DN 135:72830
 TI Phosphatidylinositol 3-kinase signaling inhibits **DAF-16**
 DNA binding and function via 14-3-3-dependent and 14-3-3-independent
 pathways
 SO Journal of Biological Chemistry (2001), 276(16), 13402-13410
 CODEN: JBCHA3; ISSN: 0021-9258
 AU Cahill, Catherine M.; Tzivion, Guri; Nasrin, Nargis; Ogg, Scott; Dore,
 Justin; **Ruvkun, Gary**; Alexander-Bridges, Maria
 AB In *Caenorhabditis elegans*, an insulin-like signaling pathway to
 phosphatidylinositol 3-kinase (PI 3-kinase) and AKT neg. regulates the
 activity of **DAF-16**, a Forkhead transcription factor.
 We show that in mammalian cells, *C. elegans* **DAF-16** is
 a direct target of AKT and that AKT phosphorylation generates 14-3-3
 binding sites and regulates the nuclear/cytoplasmic distribution of
DAF-16 as previously shown for its mammalian homologs
 FKHL and FKHL1. In vitro, interaction of AKT-phosphorylated **DAF**
-16 with 14-3-3 prevents **DAF-16** binding to
 its target site in the insulin-like growth factor binding protein-1 gene,
 the insulin response element. In HepG2 cells, insulin signaling to PI
 3-kinase/AKT inhibits the ability of a GAL4 DNA binding domain/**DAF**
-16 fusion protein to activate transcription via the
 insulin-like growth factor binding protein-1-insulin response element, but
 not the GAL4 DNA binding site, which suggests that insulin inhibits the
 interaction of **DAF-16** with its cognate DNA site.
 Elimination of the **DAF-16**/14-3-3 assocn. by mutation

of the AKT/14-3-3 sites in **DAF-16**, prevents 14-3-3 inhibition of **DAF-16** DNA binding and insulin inhibition of **DAF-16** function. Similarly, inhibition of the **DAF-16**/14-3-3 assocn. by exposure of cells to the PI 3-kinase inhibitor LY294002, enhances **DAF-16** DNA binding and transcription activity. Surprisingly constitutively nuclear **DAF-16** mutants that lack AKT/14-3-3 binding sites also show enhanced DNA binding and transcription activity in response to LY294002, pointing to a 14-3-3-independent mode of regulation. Thus, our results demonstrate at least two mechanisms, one 14-3-3-dependent and the other 14-3-3-independent, whereby PI 3-kinase signaling regulates **DAF-16** DNA binding and transcription function.

(FILE 'HOME' ENTERED AT 11:18:16 ON 16 APR 2003)

FILE 'MEDLINE, AGRICOLA, CANCERLIT, SCISEARCH, CAPLUS, MEDICONF' ENTERED
AT 11:18:50 ON 16 APR 2003

E RUVKUN G?/AU

L1 76 S E4
L2 69 DUP REM L1 (7 DUPLICATES REMOVED)
L3 11 S L2 AND DAF-16
L4 11 SORT L3 PY

=> d an ti so au ab pi l4 1-11

L4 ANSWER 1 OF 11 CAPLUS COPYRIGHT 2003 ACS
AN 1995:13340 CAPLUS
DN 122:48240
TI Daf-2, **daf-16** and daf-23: Genetically interacting
genes controlling dauer formation in *Caenorhabditis elegans*
SO Genetics (1994), 137(1), 107-20
CODEN: GENTAE; ISSN: 0016-6731
AU Gottlieb, Shoshanna; **Ruvkun, Gary**
AB Under conditions of high population d. and low food, *Caenorhabditis elegans* forms an alternative third larval stage, called the dauer stage, which is resistant to desiccation and harsh environments. Genetic anal. of some dauer constitutive (Daf-c) and dauer defective (Daf-d) mutants has revealed a complex pathway that is likely to function in particular neurons and/or responding tissues. Here the authors analyze the genetic interactions between 3 genes which comprise a branch of the dauer formation pathway that acts in parallel to or downstream of the other branches of the pathway, the Daf-c genes daf-2 and daf-23 and the Daf-d gene **daf-16**. Unlike mutations in other Daf-c genes, mutations in both daf-2 and daf-23 cause non-conditional arrest at the dauer stage. The authors epistasis anal. suggests that daf-2 and daf-23 are functioning at a similar point in the dauer pathway. First, mutations in daf-2 and daf-23 are epistatic to mutations in the same set of Daf-d genes. Second, daf-2 and daf-23 mutants are suppressed by mutations in **daf-16**. Mutations in **daf-16** do not suppress any of the other Daf-c mutants as efficiently as they suppress daf-2 and daf-23 mutants. Third, double mutants between either daf-2 or daf-23 and several other daf-d mutants exhibit an unusual interaction. Based on these results, the authors present a model for the function of daf-2, daf-23 and **daf-16** in dauer formation.

L Number	Hits	Search Text	DB	Time stamp
1	10	Ruvkun NEAR Gary	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/04/16 11:06
7	12	daf-16	USPAT; US-PGPUB; EPO; JPO; DERWENT	2003/04/16 11:06
13	11	(US-6225120-\$ or US-6472515-\$ or US-6319708-\$ or US-6135942-\$).did. or (US-20010016332-\$ or US-20010029617-\$ or US-20020037585-\$ or US-20030036079-\$).did. or (WO-9805761-\$ or WO-9851351-\$).did. or (WO-200118549-\$).did.	USPAT; US-PGPUB; EPO; DERWENT	2003/04/16 11:14

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/12, 15/63, 1/21, C07H 21/04		A1	(11) International Publication Number: WO 98/05761
			(43) International Publication Date: 12 February 1998 (12.02.98)
(21) International Application Number: PCT/US97/13914		(81) Designated States: CA, CN, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 7 August 1997 (07.08.97)			
(30) Priority Data: 60/023,382 7 August 1996 (07.08.96) US		Published With international search report.	
(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US).			
(72) Inventors: RUVKUN, Gary; 120 Herrick Road, Newton, MA 02159 (US). MORRIS, Jason; Box 574, Mather House #427, Harvard University, Cambridge, MA 02138 (US). TISSENBAUM, Heidi; 43 Chandler Street, Belmont, MA 02178 (US).			
(74) Agent: ELBING, Karen; Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110-2214 (US).			

(54) Title: AGE-1 POLYPEPTIDES AND RELATED MOLECULES AND METHODS

(57) Abstract

Disclosed are substantially pure AGE-1 polypeptides and purified DNAs, vectors, and cells encoding those polypeptides. Also disclosed are methods for determining longevity and isolating antagonists using the AGE-1 sequence.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number
WO 01/18549 A1

- (51) International Patent Classification⁷: **G01N 33/68**
- (21) International Application Number: **PCT/US00/24487**
- (22) International Filing Date:
7 September 2000 (07.09.2000)
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/152,825 7 September 1999 (07.09.1999) **US**
- (71) Applicant (for all designated States except US): **NEURO-GENETICS, INC.** [US/US]; 4310 La Jolla Village Drive #400, San Diego, CA 92122 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HENDERSON, Samuel, T.** [US/US]; 2110 West 10th Avenue #203, Broomfield, CO 80020 (US). **JOHNSON, Thomas, E.** [US/US]; 1121 West Enclave Circle, Louisville, CO 80027 (US).
- (74) Agents: **SWANSON, Barry, J. et al.**; Swanson & Bratschun, L.L.C., Suite 330, 1745 Shea Center Drive, Highlands Ranch, CO 80129 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/18549 A1

(54) Title: **THERAPIES AND REAGENTS FOR INCREASING STRESS RESISTANCE AND LIFE SPAN**

(57) Abstract: The invention comprises methods for identifying agents that can activate *C. elegans* DAF-16 and human homologs thereof, or their corresponding genes, whereby cytoprotective effects in cells may be induced. Such cytoprotective effects can result in enhanced environmental stress resistance, increased life span and improved late life vigor without significant inhibition of insulin-signaling pathways. The invention also comprises the therapeutic agents identified and methods of treatment using the agents.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/24487

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 51351 A (GEN HOSPITAL CORP) 19 November 1998 (1998-11-19) cited in the application	1-51
Y	the whole document	1,5,6, 10,16, 20,28, 32,47,51
Y	RENA G ET AL: "Phosphorylation of the transcription factor forkhead family member FKHR b protein kinase B." JOURNAL OF BIOLOGICAL CHEMISTRY, (1999 JUN 11) 274 (24) 17179-83. , XP000981982 the whole document	1,5,6, 10,16, 20,28, 32,47,51
	--- -/-- ---	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

8 February 2001

Date of mailing of the international search report

22/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Gundlach, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/24487

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KOPS G J ET AL: "Direct control of the Forkhead transcription factor AFX by protein kinase B." NATURE, (1999 APR 15) 398 (6728) 630-4. , XP000955388 the whole document -----	1,5,6, 10,16, 20,28, 32,47,51

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/24487

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9851351 A	19-11-1998	AU 7494198 A	08-12-1998
		EP 1019092 A	19-07-2000
		PL 336858 A	17-07-2000
		HU 0002199 A	28-09-2000
<hr/>			